

PRODUCTIVITY (ABOVE- AND BELOWGROUND) OF *FESTUCA ARUNDINACEA* SCHREB FOLLOWING DIFFERENT DEFOLIATIONS

Hennie Snyman

Paul Malan



August 2, 2016

T: 051 401 9111 info@ufs.ac.za www.ufs.ac.za

© Copyright reserved
Kopiereg voorbehou

UNIVERSITY OF THE
FREE STATE
UNIVERSITEIT VAN DIE
VRYSTAAT
YUNIVESITHI YA
FREISTATA




UFS·UV
NATURAL AND
AGRICULTURAL SCIENCES
NATUUR- EN
LANDBOUWETENSAPPE


GREENHOUSE TRIAL – DEFOLIATION



RESULTS (ABOVEGROUND PHYTOMASS PRODUCTION)



RESULTS (ABOVE- AND BELOWGROUND PHYTOMASS PRODUCTION)









CONCLUSION



- Longer resting (recovery) periods are essential for healthy root systems – plant survival

Thank You
Dankie



T: 051 401 9111 info@ufs.ac.za www.ufs.ac.za

© Copyright reserved
Kopiereg voorbehou

UNIVERSITY OF THE
FREE STATE
UNIVERSITEIT VAN DIE
VRYSTAAT
YUNIVESITHI YA
FREISTATA



UFS·UV
NATURAL AND
AGRICULTURAL SCIENCES
NATUUR- EN
LANDBOUWETENSAPPE